

Largeness of LERF and 1-relator groups

J. O. Button
 Selwyn College
 University of Cambridge
 Cambridge CB3 9DQ
 U.K.
 jb128@dpmms.cam.ac.uk

Abstract

We consider largeness of groups given by a presentation of deficiency 1, where the group is respectively free-by-cyclic, LERF or 1-relator. We give the first examples of (finitely generated free)-by- \mathbb{Z} word hyperbolic groups which are large, show that a LERF deficiency 1 group with first Betti number at least 2 is large or $\mathbb{Z} \times \mathbb{Z}$ and show that 2-generator 1-relator groups where the relator has height 1 obey the dichotomy that either the group is large or all its finite images are metacyclic.

1 Introduction

A finitely generated group G is said to be large if it has a finite index subgroup possessing a homomorphism onto a non-abelian free group. This has a range of implications: for instance G is SQ-universal (which means that every countable group is a subgroup of a quotient of G), G has uniformly exponential word growth, G has the largest possible subgroup growth for finitely generated groups (which is of strict type n^n) and G has infinite virtual first Betti number. If we restrict ourselves to finitely presented groups and define the deficiency of a finite presentation to be the number of generators minus the number of relators, a well known result of B. Baumslag and S. J. Pride in [5] is that groups with a presentation of deficiency at least 2 are large.

The results in this paper follow on from [24] where the question considered was which groups with a deficiency 1 presentation are large. Clearly \mathbb{Z} and $\mathbb{Z} \times \mathbb{Z}$ are not, and neither are the soluble Baumslag-Solitar groups $BS(1, n)$, where the Baumslag-Solitar group $BS(m, n)$ has the presentation $\langle x, y | yx^my^{-1} = x^n \rangle$. There are other cases, such as $BS(2, 3)$, but no more residually finite examples are known (indeed all others are “far from being residually finite” in a sense that will be made precise in Section 4).

However in [24] many families of deficiency 1 groups which are all large were found. In particular if F_n is the free group of rank n then it was shown that free-by-cyclic groups $F_n \rtimes_{\alpha} \mathbb{Z}$ are large for $n \geq 2$ if they contain $\mathbb{Z} \times \mathbb{Z}$, which is equivalent here to not being a word hyperbolic group. The question of whether $F_n \rtimes_{\alpha} \mathbb{Z}$ is large in the word hyperbolic case was left open, and up until now not a single large example was known. In Section 2 we show that if α is a reducible automorphism then $F_n \rtimes_{\alpha} \mathbb{Z}$ is large provided that free-by-cyclic groups have finite index subgroups with first Betti number at least 2. This was raised by A. Casson in [6] Question 12.16. Although still unknown in the word hyperbolic case, if α is a specific reducible automorphism then we merely require for largeness that two particular free-by-cyclic groups have virtual first Betti number at least 2: these are the one obtained by restricting (a suitable power of) α to the invariant free factor, and the free-by-cyclic group formed by quotienting out the invariant free factor. By taking a specific word hyperbolic group in the literature which is of the form $F_3 \rtimes_{\alpha} \mathbb{Z}$ and using it to make a reducible automorphism, we obtain Corollary 2.4 which gives the first group of the form $F_n \rtimes_{\alpha} \mathbb{Z}$ which is known to be both word hyperbolic and large. It double covers a group with the same properties which has a most succinct presentation: $\langle t, a | t^6 a t^{-4} a^{-1} t^{-2} a^{-1} \rangle$.

We have mentioned that the property of residual finiteness should increase the chances of a deficiency 1 presentation being large. For instance all groups of the form $F_n \rtimes_{\alpha} \mathbb{Z}$ are residually finite. In Section 3 we look at deficiency 1 groups which are LERF (also known as subgroup separable). This is considerably stronger than residual finiteness so we would expect these groups to be large (with three obvious exceptions). Once again though the problem is finding a finite cover with first Betti number at least 2. We prove in Theorem 3.3 that if G is a LERF group with a presentation of deficiency 1 and has such a finite cover then G is large or the fundamental group of the torus or Klein bottle. A recent result in [41] of D. Kochloukova is used in Theorem 3.1 to show that the only possible exceptions to LERF deficiency 1 groups being large (apart from \mathbb{Z} and these two groups) are word hyperbolic

groups of the form $F_n \rtimes_\alpha \mathbb{Z}$. This at least gives us in Corollary 3.2 that all but these three deficiency 1 LERF groups are SQ-universal, but we are prevented from concluding largeness until Casson's question is settled (at least in the LERF case but even here this seems open). We also look at the question of a type of Tits alternative for deficiency 1 groups which would say that such a group is either soluble or contains a non-abelian free group. The result in [41] mentioned above very nearly established this but one case is still to be resolved. We show in Corollary 3.4 that this is true if G has a finite index subgroup with first Betti number at least 2. However it is not true that all finitely generated subgroups of G will either be virtually soluble or contain a non-abelian free group, and Example 3.5 is such a group which has deficiency exactly 1.

Another much studied class of groups are those with a 1-relator presentation. The intersection of 1-relator and deficiency 1 groups is the class of 2-generator 1-relator presentations. Although groups of this form have strong properties, it is not known which ones are large. Moreover we would like to be able to deduce largeness using only information obtained directly from the presentation rather than needing to know a priori that the group has special properties such as residual finiteness.

In Section 4 we settle this question for a particular class of 2-generator 1-relator presentations. Given any relator in 2 variables we can make a change of basis of F_2 to $\{a, t\}$ such that the exponent sum of t in the relator is zero. We say that r has height 1 if appearances of $t^{\pm 1}$ in r are such that t alternates with t^{-1} . Although this is a restricted set of relators, it is the case that nearly all of the 2-generator 1-relator groups in the literature which have unusual or nasty properties (we review these in that section) are given by height 1 words.

We establish a major dichotomy of groups G with height 1 presentations in Theorem 4.1 which states that either G is large or all its finite images are metabelian. If G is not a soluble Baumslag-Solitar group but is in the latter case then G contains a non-abelian free group, so is far from being metabelian and hence far from being residually finite. We also apply a famous result of Zelmanov on pro- p groups which allows us in Corollary 4.2 to distinguish between the two cases: G is large if and only if it has a finite index subgroup whose abelianisation requires at least 3 generators, which is a condition that can easily be checked on a computer.

It is not true that a large height 1 group is necessarily residually finite. However in Corollary 4.6 we give the first example of a 2-generator 1-relator

presentation of a group G where $\beta_1(G) = 2$ and the only finite images of G are abelian but G is not equal to $\mathbb{Z} \times \mathbb{Z}$. This means that it is not enough to conclude largeness (or equality with $\mathbb{Z} \times \mathbb{Z}$) for a deficiency 1 group with first Betti number at least 2 (even in the 1-relator case). However this is unknown if the first Betti number is at least 3, or even if the abelianisation of the group requires at least 3 generators.

Section 5 is a collection of open questions encountered during the preparation of this work which, although some of these might be well known, we have not found in the standard problem lists. They are all on finitely generated and finitely presented groups and the list begins with the most wide ranging questions and then gradually specialises, ending with unsolved problems that are the most relevant to this paper.

The author would like to thank J. Hillman for bringing his attention to the paper [41].

2 Reducible Free-by-Cyclic Groups

A powerful method for proving largeness directly from a given finite presentation is to apply Howie's result which is Theorem A in [39]. This tells us that if there is a homomorphism χ from a finitely presented group G onto \mathbb{Z} such that the Alexander polynomial $\Delta_{G,\chi}(t) \in \mathbb{Z}[t^{\pm 1}]$ relative to χ is identically zero then G is large (indeed the proof shows that a finite index subgroup containing $\ker \chi$ surjects onto a non-abelian free group). However not only do we need the abelianisation $\overline{G} = G/G'$ of G to be infinite in order to have a homomorphism onto \mathbb{Z} in the first place, we also have that if $\overline{G} = \mathbb{Z} \times T$ for T finite then $|\Delta_{G,\chi}(1)|$ is the order of T . Thus we will not be able to use this criterion for largeness unless the first Betti number $\beta_1(G)$ is at least two, or we can find a finite index subgroup with this property. (In fact Howie's theorem is also true if the mod p Alexander polynomial $\Delta_{G,\chi}^p \in \mathbb{F}[t^{\pm 1}]$ for $\mathbb{F} = \mathbb{Z}/p\mathbb{Z}$ is zero and we will use this in later sections, but for now we will stick to the characteristic zero version.)

Given a finitely presented group $G = \langle x_1, \dots, x_m | r_1, \dots, r_n \rangle$ which does have a homomorphism χ onto \mathbb{Z} where $K = \ker \chi$, we can regard K/K' as a $\mathbb{Z}[t^{\pm 1}]$ -module where t acts by conjugation on K using an element of $\chi^{-1}(1)$. Moreover we can obtain a finite presentation for this module K/K' by using the Reidemeister-Schreier rewriting process to go from a group presentation for G to a group presentation for K , and then abelianising the relations.

Although this will result in infinitely many group relations there are only finitely many orbits under the action of t . The result is an $(m - 1)$ by n presentation matrix M for K/K' and we can assume that $n \geq m - 1$ by adding zero columns if necessary. We then define the Alexander polynomial $\Delta_{G,\chi} \in \mathbb{Z}[t^{\pm 1}]$ to be the highest common factor of the $(m - 1)$ by $(m - 1)$ minors of M ; it is the same (up to units in $\mathbb{Z}[t^{\pm 1}]$) for any finite presentation of G .

The fact that a zero Alexander polynomial implies largeness is particularly useful for groups G with a presentation of deficiency 1 because the resulting matrix is square and so we are merely evaluating the determinant to obtain the Alexander polynomial. In particular if we find that one row or column consists entirely of zeros then we immediately conclude largeness. We do have the problem mentioned above that we need $\beta_1(G) \geq 2$ for this to happen, however another advantage of deficiency 1 presentations is that for any finite index subgroup H of G (for which we write $H \leq_f G$) the Reidemeister-Schreier rewriting process results in a deficiency 1 presentation for H . As H is large if and only if G is, we can hope that there is a subgroup H with $\beta_1(H) \geq 2$.

A large class of deficiency 1 presentations come from the free-by-cyclic groups: let F be a free group and α an automorphism of F . Then we can form the semidirect product (also called the mapping torus) $F \rtimes_{\alpha} \mathbb{Z}$. If F_n is the free group of rank n with free basis x_1, \dots, x_n then $F_n \rtimes_{\alpha} \mathbb{Z}$ has the presentation

$$\langle x_1, \dots, x_n, t | tx_1t^{-1} = \alpha(x_1), \dots, tx_nt^{-1} = \alpha(x_n) \rangle. \quad (1)$$

The following facts are known about free-by-cyclic groups $F \rtimes_{\alpha} \mathbb{Z}$; see [24] Section 5 and references within.

- (1) If F is of infinite rank then $F \rtimes_{\alpha} \mathbb{Z}$ may be finitely or infinitely generated. If $F \rtimes_{\alpha} \mathbb{Z}$ is infinitely generated then it need not be large, nor residually finite, but if it is finitely generated then it is residually finite and finitely presented. Moreover it has a presentation with deficiency at least 2, so is large.
- (2) $F_n \rtimes_{\alpha} \mathbb{Z}$ is residually finite, has deficiency exactly equal to 1 and any finite index subgroup is also of the form $F_m \rtimes_{\beta} \mathbb{Z}$ so also has deficiency exactly 1.
- (3) $F_n \rtimes_{\alpha} \mathbb{Z}$ is word hyperbolic precisely when it does not contain a subgroup isomorphic to $\mathbb{Z} \times \mathbb{Z}$. If it does and $n \geq 2$ then it is large. However it is not known whether word hyperbolic groups of the form $F_n \rtimes_{\alpha} \mathbb{Z}$ are large. A problem in [6] due to Casson is whether a group $G = F_n \rtimes_{\alpha} \mathbb{Z}$ with $n \geq 2$

always has $H \leq_f G$ with $\beta_1(H) \geq 2$. Whilst this is true if G contains $\mathbb{Z} \times \mathbb{Z}$, it is unknown in general if G is word hyperbolic and so we will need to take this as an assumption.

An automorphism α of the free group F_n for $n \geq 2$ is said to be reducible if there exist proper non-trivial free factors R_1, \dots, R_k of F_n such that the conjugacy classes of R_1, \dots, R_k are permuted transitively by α (see [9]).

Theorem 2.1 *Assume that any group of the form $F_n \rtimes_\alpha \mathbb{Z}$ for $n \geq 2$ has a finite index subgroup with first Betti number at least 2. Then if $G = F_n \rtimes_\alpha \mathbb{Z}$ for α a reducible automorphism, we have that G is large.*

Proof. Our group G will have a presentation as in (1) and so for each positive integer k there exists the cyclic cover of G which is the index k subgroup $F_n \rtimes_{\alpha^k} \mathbb{Z}$ generated by t^k and F_n . If α is reducible then on replacing α^k by α we can assume $F_n = A * B$ with A and B proper free factors and $\alpha(A)$ is sent to a conjugate of A . However on now composing α with an appropriate inner automorphism (which does not change the free-by-cyclic group) we can assume that $\alpha(A) = A$. From now on this will be our G .

We define the free-by-cyclic group G_r by restricting α to A . Our assumption means that we can take a finite index subgroup of G_r with first Betti number at least 2 (and note that if A has rank 1 then we can do this as well). But every finite index subgroup of $F_n \rtimes_\alpha \mathbb{Z}$ contains one of the form $F_m \rtimes_{\alpha^k} \mathbb{Z}$ where $F_m \leq_f F_n$ and $\alpha^k(F_m) = F_m$. Moreover the first Betti number does not decrease in finite covers. Doing this for G_r , we have $A_0 \leq_f A$ and a power of α fixing A_0 (which we again replace by α) to get a finite index subgroup $L = \langle t, A_0 \rangle$ of G_r with $\beta_1(L) \geq 2$. However a result of Marshall Hall Jnr. states that as A_0 is a finitely generated subgroup of $F_n = A * B$, there is a finite index subgroup E of F_n with A_0 a free factor, so we have $E = A_0 * S$ for some S and, as A_0 does not have finite index in F_n , S is non trivial. Again taking a power of t , we can assume that $\alpha^j(E) = E$ as there are only finitely many subgroups in F_n of each finite index.

We are now ready to look at a finite presentation for the finite index subgroup $J = \langle t^j, E \rangle$ of G . On taking a free basis a_1, \dots, a_l for A_0 and a similar one for S , we find that the first l relations in the deficiency 1 presentation for L as in (1) are of the form $ta_i t^{-1} = w_i(a_1, \dots, a_l)$ where w_i are reduced words in F_l . However $\beta_1(L)$ being at least 2 means that we can change the free basis for A_0 so that the first relation is actually $ta_1 t^{-1} = a_1 c(a_1, \dots, a_l)$ where c is a reduced word in the commutator subgroup F_l'

of F_l . Now suppose there exists a surjective homomorphism χ from J to \mathbb{Z} with the property that t and all of A_0 are in $\ker \chi$. When we form the square matrix M with entries in $\mathbb{Z}[x^{\pm 1}]$ in order to calculate the polynomial $\Delta_{J,\chi}(x)$, all the letters appearing in the first relation are in the kernel of χ . Therefore we have that the first column of M consists in turn of the exponent sum of t, a_1, \dots, a_l and then zeros in the other places because the latter rows correspond to generators in this presentation for J which do not even appear in the first relation. However the exponent sums of the generators t, a_1, \dots, a_l that do appear are all zero so we have a zero column in a square matrix, meaning that $\Delta_{J,\chi}$ is the zero polynomial so J , and G , are large.

In order to find such a homomorphism, we must again use our assumption on finite covers with first Betti number at least 2. From G we obtained the “reduced” free-by-cyclic group G_r by restricting α to A . Although the definition of a reducible automorphism means that we cannot assume $\alpha(B) = B$ even if $\alpha(A) = A$, we can form the “quotient” free-by-cyclic group G_q by taking the “quotient” automorphism $\tilde{\alpha}$ of B . This is formed by letting $\pi : A * B \rightarrow B$ be the homomorphism with kernel the normal closure of A and then we define $\tilde{\alpha}(b) = \pi\alpha(b)$ for $b \in B$. Note that $\tilde{\alpha}$ is surjective because α and π are, and the Hopfian property of finitely generated free groups means that $\tilde{\alpha}$ is an automorphism. Also there is a natural homomorphism θ from G to $G_q = \langle s, B \rangle$ given by sending t to s and “ignoring” A ; this is well defined because $\alpha(A) = A$.

Now we apply our assumption to obtain a finite index subgroup H of G_q with $\beta_1(H) \geq 2$ and without loss of generality we can assume H is of the form $\langle s^i, C \rangle$ for $C \leq_f B$ and some $i \in \mathbb{N}$. In fact as cyclic covers of H will also have first Betti number at least 2, we can replace i with j by taking multiples so that they are both equal to ij . As $\beta_1(H) \geq 2$, we must have a homomorphism $\tilde{\chi}$ from H onto \mathbb{Z} with $\tilde{\chi}(s^j) = 0$. We then consider the subgroup of G which is $\theta^{-1}(H) \cap J$. As each of these subgroups has finite index in G , so does their intersection. We can take our finite presentation above for J and use this to rewrite for the finite index subgroup $\theta^{-1}(H) \cap J$. Note that t^j is in $\theta^{-1}(H)$ and this presentation will also have deficiency 1. Also the generators that we used for A_0 are in $\ker \theta$ so are in $\theta^{-1}(H)$ too. Hence in going from our old presentation to our new one, our valuable relation survives intact. But the homomorphism $\tilde{\chi}\theta : \theta^{-1}(H) \rightarrow \mathbb{Z}$ can be restricted to $\theta^{-1}(H) \cap J$ which has finite index in $\theta^{-1}(H)$. Without loss of generality this restriction is surjective and moreover all the generators in our special relation end up being sent to the identity by $\tilde{\chi}\theta$ so we are done.

□

Corollary 2.2 *Assume that any group $F_n \rtimes_\alpha \mathbb{Z}$ for $n \geq 2$ has a finite index subgroup with first Betti number at least two. Suppose that the group G can be written in the form $F_n \rtimes_\alpha \mathbb{Z}$ and there exists a finitely generated subgroup A of F_n which is non-trivial and of infinite index in F_n , whose conjugacy class has a finite orbit under α . Then G is large.*

Proof. By taking cyclic covers and an inner automorphism, we can assume that $\alpha(A) = A$. Although A may not be a free factor of F_n , we can again use M. Hall Jr.'s result to find S (with A and S non-trivial) such that $A * S = F_m$ and F_m has finite index in F_n . We now take the appropriate power α^i of α that fixes F_m . Then the subgroup $H = F_m \rtimes_{\alpha^i} \mathbb{Z}$ has finite index in G , with α^i reducible when restricted to F_m , so H is large by Theorem 2.1.

□

These results are all very well but we have not seen a single example of a word hyperbolic group of the form $F_n \rtimes_\alpha \mathbb{Z}$ which is large. Although it seems as if we need to wait for Casson's question on finite index subgroups with first Betti number at least two to be settled positively, we can manage without this in specific cases. If we have an automorphism α of $F_n = A * B$ with $\alpha(A) = A$, where A and B are proper free factors, and we form the free-by-cyclic group G then the proof of Theorem 2.1 was set out so that it is enough to find finite index subgroups with first Betti number 2 of both the restriction $G_r = A \rtimes_\alpha \mathbb{Z}$ and the quotient $G_q = B \rtimes_{\bar{\alpha}} \mathbb{Z}$. If we have a presentation for G as in (1) and our free basis for F_n is obtained by putting together ones for A and B then we instantly get finite presentations for G_r and G_q which we can then feed into a computer and ask it to enumerate finite index subgroups and their abelianisations.

This allows us to check largeness of free-by-cyclic groups G formed by reducible automorphisms α . If we require a word hyperbolic example then it is necessary that G contains no $\mathbb{Z} \times \mathbb{Z}$ subgroup, or equivalently α has no periodic conjugacy classes. Moreover this is sufficient for a group of the form $F_n \rtimes \mathbb{Z}$ to be word hyperbolic by [7], [8] and [16]. Thus it is straightforward to create reducible word hyperbolic examples by a “doubling” process.

Lemma 2.3 *If $A \rtimes_\alpha \mathbb{Z}$ is word hyperbolic where A is isomorphic to F_n and α is an automorphism of A then $(A * B) \rtimes_\alpha \mathbb{Z}$ is also word hyperbolic, where B is a copy of A and the action of α on B is the same as on A .*

Proof. Suppose we have $w \in A * B$ with $\alpha^k(w)$ equal to a conjugate of w for $k \geq 1$. Then w is not in A or B so we can ensure (by conjugation in $A * B$ if necessary) that $w = a_1 b_1 \dots a_n b_n$ where $a_i \in A - \{e\}$ and $b_i \in B - \{e\}$. Now on setting $a'_i = \alpha^k(a_i)$ which is in A and $b'_i = \alpha^k(b_i)$ in B , we have that w and $a'_1 b'_1 \dots a'_n b'_n$ are conjugate, with both words cyclically reduced in a free group, meaning that there must be j so that (taking subscripts modulo n) $a'_i = a_{i+j}$ and $b'_i = b_{i+j}$. But then $\alpha^{kn}(a_i) = (a_i)$, giving a $\mathbb{Z} \times \mathbb{Z}$ subgroup in $A \rtimes_\alpha \mathbb{Z}$. \square

Note that in this case if $G = (A * B) \rtimes_\alpha \mathbb{Z}$ then both G_r and G_q are isomorphic to the original group $A \rtimes_\alpha \mathbb{Z}$.

We now need to find explicit examples of word hyperbolic groups of the form $F_n \rtimes_\alpha \mathbb{Z}$. To achieve this we use results in [34] and [56]. An automorphism α of F_n gives rise to an automorphism of the abelianisation \mathbb{Z}^n of F_n which is well defined for outer automorphisms, as is the definition of reducibility of α . In the former paper Corollary 2.6 states that if we have an outer automorphism O of F_n such that the characteristic polynomial of the induced automorphism of \mathbb{Z}^n is a PV-polynomial (which means that the polynomial is monic and has exactly one root with modulus greater than one (counted with multiplicity) and no roots on the unit circle) then O^k is irreducible for $k \geq 1$. They then obtain as Corollary 2.8 that if α is an automorphism of F_n for $n \geq 3$ where the corresponding characteristic polynomial is a PV-polynomial then $\alpha(w) = w$ implies that $w = e$. Moreover α can have no periodic conjugacy classes because α^k will also have the same property, and if α (or α^k) sends w to a conjugate then we multiply α (or α^k) by an inner automorphism which does not change the characteristic polynomial.

The latter Corollary is proved by applying [9] Theorem 4.1 which states that if O^k is irreducible for $k \geq 1$ and O fixes a conjugacy class then O is geometrically realised by a pseudo-Anosov homeomorphism of a compact surface with one boundary component. But [56] shows that automorphisms of F_n with a characteristic polynomial that is a PV-polynomial are not geometrically realisable for $n \geq 3$ by considering the eigenvalues (note that $F_2 \rtimes_\alpha \mathbb{Z}$ is never word hyperbolic because the commutator of the generators gives rise to a conjugacy class that is fixed or of period 2). An example is given: $G_0 = F_3 \rtimes_\alpha \mathbb{Z}$ where $\alpha(x) = y, \alpha(y) = z, \alpha(z) = xy$.

Corollary 2.4 *The F_6 -by- \mathbb{Z} group*

$$G = \langle t, a, b, c, x, y, z \mid tat^{-1} = b, tbt^{-1} = c, tct^{-1} = ab, \dots \rangle$$

$$txt^{-1} = y, tyt^{-1} = z, tzt^{-1} = xy$$

is word hyperbolic and large.

Proof. That G is word hyperbolic follows from the facts quoted above: namely $G_0 = F_3 \rtimes_{\alpha} \mathbb{Z}$ has no periodic conjugacy classes so Lemma 2.3 shows that G also has no periodic conjugacy classes thus is word hyperbolic. That G is large follows from applying the proof of Theorem 2.1 to G with $G_r = G_q = G_0$, and from the output of a computer. Inputting the presentation of G_0 into MAGMA and asking for the abelianisation of its low index subgroups, we find (after a bit of a wait, although it is more quickly checked) that G_0 has an index 14 subgroup (with generators $x, y, z^2, zxz^{-1}, zyz^{-1}, zt^{-7}$) with abelianisation $C_2 \times C_4 \times \mathbb{Z} \times \mathbb{Z}$. □

We also note that G is the double cover of the automorphism which sends in turn a to x to b to y to c to z to ab . Here all other generators except t and a can be eliminated to get the 2-generator 1-relator F_6 -by- \mathbb{Z} word hyperbolic large group $\langle t, a | t^6 a t^{-4} a^{-1} t^{-2} a^{-1} \rangle$.

3 LERF groups of deficiency 1

Not all groups with a deficiency 1 presentation can be large, as evidenced by the Baumslag-Solitar groups $BS(m, n) = \langle a, t | ta^m t^{-1} = a^n \rangle$ for m, n non-zero integers, where we can take without loss of generality $m > 0$ and $|n| \geq m$. We have that $BS(m, n)$ is large if and only if m and n are not coprime. For $m = 1$ we have a soluble group (and it is known that a virtually soluble group of deficiency 1 must be isomorphic to $BS(1, n)$ or to \mathbb{Z}) but otherwise $BS(m, n)$ contains a non-abelian subgroup, thus it is not true that Baumslag-Solitar groups are either virtually soluble or large. However most of these groups are not residually finite (for this we require $m = 1$ or $m = |n|$) and so if we stick to residually finite Baumslag-Solitar groups, we do have this dichotomy. Thus we could consider residually finite groups of deficiency 1: in fact we know of no example of such a group which is not virtually soluble but not large.

In order to make progress we impose an even tighter condition on our deficiency 1 group G , which is that it is LERF (locally extended residually finite, also known as subgroup separable). This means that every finitely generated subgroup is an intersection of finite index subgroups. The big

advantage of this is a result of Lubotzky in [45] that if a LERF group G can be written as an HNN extension $H*_\phi$ which is non-ascending, that is both the domain A of ϕ and the image B are strictly contained in H , then G is large if A is finitely generated. This is because G surjects to an HNN extension of a finite group which is virtually free, and the LERF property applied to A means that this HNN extension will also be non-ascending, thus it is virtually free but not virtually cyclic.

Theorem 3.1 *If G is LERF and of deficiency 1 then either G is large or G is of the form $F_n \rtimes \mathbb{Z}$.*

Proof. As there must be a surjective homomorphism χ from G to \mathbb{Z} , we have by [12] that the finitely presented group G is an HNN extension of a finitely generated group H with finitely generated associated subgroups A and B . We then have that if this HNN extension is not ascending then G is large by the above. But if say $H = A$ then $G = \langle t, H \rangle$ where t is the stable letter and we have $tHt^{-1} \subseteq H$. By a result of Blass and P. M. Neumann, a LERF group cannot have a finitely generated subgroup conjugate to a proper subgroup of itself so $tHt^{-1} = H$ and we conclude that $H \trianglelefteq G$ with $G = H \rtimes \mathbb{Z}$. Now we can use a recent result of Kochloukova. We have by [36] Theorem 6 that if G has deficiency 1 and is an ascending HNN extension of a finitely generated group then G has geometric (hence cohomological) dimension at most 2. But Theorem 3 in [41] is as follows: Let G be a non-trivial group with a finite $K(G,1)$ CW-complex of dimension n with Euler characteristic 0. Suppose that N is a normal subgroup of G containing G' which is of homological type FP_{n-1} and G/N is cyclic-by-finite. Then N is of type FP_n . Consequently we conclude by putting H equal to N that H is of type FP_2 , but a result of Bieri gives us that the cohomological dimension of H is 1 and thus H is free. \square

It should be noted that groups of the form $F_n \rtimes \mathbb{Z}$ are not necessarily LERF.

Consequently if we had that $F_n \rtimes \mathbb{Z}$ was large for $n \geq 2$ we would have the strongest possible result: either a deficiency 1 LERF group is large or it is \mathbb{Z} , $\mathbb{Z} \times \mathbb{Z}$ or the Klein bottle group $BS(1, -1)$. However we can move in two directions from this result. The first is to relax the property of largeness to that of G being SQ-universal (every countable group is a subgroup of a quotient of G) which in turn is stronger than containing the free group F_2 . As groups of the form $F_n \rtimes \mathbb{Z}$ for $n \geq 2$ are either large by [24] or hyperbolic, hence SQ-universal by [53], we immediately get:

Corollary 3.2 *A LERF group of deficiency 1 is either SQ-universal or is one of \mathbb{Z} , $\mathbb{Z} \times \mathbb{Z}$, $BS(1, -1)$.*

A conjecture of P. M. Neumann from 1973 is that there is a dichotomy for any 1-relator group: either it is SQ-universal or it is a soluble Baumslag-Solitar group or \mathbb{Z} . Of course if there are at least 3 generators then later it was shown that we have largeness but the conjecture remains open in the 2-generator 1-relator case. Corollary 7.5 in [24] showed this to be true if the group is LERF, so in light of Corollary 3.2 we wonder if P. M. Neumann's conjecture extends to all deficiency 1 groups, or (more cautiously) to all residually finite deficiency 1 groups.

The other improvement that we can make to Theorem 3.1 is to give further conditions guaranteeing largeness. It might not be a surprise in light of Section 2 that once again it comes down to requiring first Betti number at least two.

Theorem 3.3 *If G is LERF, of deficiency 1 and has a finite index subgroup with first Betti number at least 2 then G is large or $\mathbb{Z} \times \mathbb{Z}$ or the Klein bottle group.*

Proof. We assume that $\beta_1(G) = b \geq 2$ and is LERF (which is a property preserved by all subgroups). We consider the BNS (Bieri-Neumann-Strebel) invariant of G which is an open subset of S^{b-1} and which gives information on the finite generation of kernels of non-trivial homomorphisms from G to \mathbb{Z} . By regarding the rationally defined points of S^{b-1} as equivalence classes of homomorphisms χ, χ' from G to \mathbb{Z} according to the relation $\chi = q\chi'$ for $q \in \mathbb{Q}$ and $q > 0$, we have that $[\chi]$ is in Σ if and only if G can be expressed as an ascending HNN extension $\langle t, H \rangle$ with associated homomorphism χ (that is $\chi(t) = 1$ and $\chi(H) = 0$) and with H finitely generated. Moreover $\ker \chi$ is finitely generated if and only if $[\chi]$ and $[-\chi]$ are both in Σ . Now for a LERF group G , if there exists χ with neither of $[\pm\chi]$ in Σ then G is large, as in Theorem 3.1. Otherwise one of $[\pm\chi]$ is in Σ for every homomorphism χ from G onto \mathbb{Z} . But as the LERF condition means that we cannot have G equal to a strictly ascending HNN extension $\langle t, H \rangle$ where H is finitely generated, we must have both $[\pm\chi] \in \Sigma$ for all possible χ .

Now we use a result of Dunfield in [26]. One can regard the (multivariable) Alexander polynomial $\Delta_G(t_1, \dots, t_b)$ as a finite set of lattice points in \mathbb{Z}^b labelled with a non-zero integer by taking the monomials in Δ_G with non-zero coefficient. We can then form the Newton polytope $N(\Delta_G) \subseteq \mathbb{R}^b$ which

is the convex hull of these points. Then [26] Theorem 5.1 states that if $D(\Delta_G)$ is the dual of $N(\Delta_G)$ in \mathbb{R}^b (so that faces of dimension i become faces of dimension $b - i - 1$) and F_1, \dots, F_k are the $(b - 1)$ -dimensional faces of $D(\Delta_G)$ whose corresponding vertices of $N(\Delta_G)$ have coefficient ± 1 then Σ is contained in the projection of the interiors of the F_i to S^{b-1} .

So far we have not used the fact that $b \geq 2$. Now when $b = 1$ we get $S^0 = \{\pm 1\}$ and the result above is saying that if χ is the unique surjective homomorphism (up to sign) from G to \mathbb{Z} then $\ker \chi$ being finitely generated implies that the highest and lowest terms of $\Delta_G(t)$ are monic. However if $b \geq 2$ then the fact that $[\chi] \in \Sigma^{b-1}$ for all homomorphisms χ implies that $\Delta_G = 1$. This is because if there are $n \geq 2$ vertices of $N(\Delta_G)$ then we have $(b - 1)$ -dimensional faces F_1, \dots, F_n of $D(\Delta_G)$, and so when we project the interiors of those faces which are obtained from the ± 1 coefficients, we do not cover all of S^{b-1} because we miss the lower dimensional faces where pairs of elements of F_1, \dots, F_n meet. Moreover there will be rationally defined points which are not covered and therefore homomorphisms χ with $[\chi]$ not in Σ .

Hence if G is LERF and has $\beta_1(G) \geq 2$ then the only way that G fails to be large is if $\Delta_G = 1$, with $\ker \chi$ being finitely generated for all homomorphisms χ onto \mathbb{Z} . However by [24] Theorem 3.1 we have for deficiency 1 groups that

$$\Delta_{G,\chi}(t) = (t - 1)\Delta_G(t_1^{n_1}, \dots, t_b^{n_b})$$

where n_i is the image under χ of any element in G which projects to t_i under the natural homomorphism from G to its free abelianisation \mathbb{Z}^b . Consequently if $\Delta_G = 1$ then for any χ we have $\Delta_{G,\chi}$ equal to $t - 1$. But the degree of $\Delta_{G,\chi}$ is the dimension of the \mathbb{Q} -vector space $H_1(\ker \chi; \mathbb{Q}) = H_1(\ker \chi; \mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{Q}$, for which we write $\beta_1(\ker \chi; \mathbb{Q})$, and from [41] Theorem 3 we have that $\ker \chi$ is free, so it must be free of rank 1 and therefore $G = \mathbb{Z} \times \mathbb{Z}$.

Finally if G has deficiency 1 and has a finite index subgroup $H = \mathbb{Z} \times \mathbb{Z}$ then G is either $\mathbb{Z} \times \mathbb{Z}$ or the Klein bottle group.

□

We will see in Corollary 4.6 that this theorem is not true if the LERF condition is removed. If it is weakened to residually finite then this is unknown.

We can also finish off the proof without needing [41] Theorem 3 by considering the Alexander polynomial $\Delta_{G,\chi}^p$ over $\mathbb{Z}/p\mathbb{Z}$. Although it is not true for a general finite presentation, for the deficiency 1 case this is just the Alexander polynomial $\Delta_{G,\chi}$ over \mathbb{Z} reduced modulo p . Consequently we must have $\beta_1(\ker \chi; \mathbb{Z}/p\mathbb{Z}) = 1$ and so $\beta_1(G; \mathbb{Z}/p\mathbb{Z}) \leq 2$. But if this is true for all finite

index subgroups H of G then $H/H' = \mathbb{Z} \times \mathbb{Z}$ which cannot happen if G is residually finite by [24] Proposition 3.4 unless $G = \mathbb{Z} \times \mathbb{Z}$.

As for attempting to apply the above techniques to deficiency 1 groups which are not LERF, we can squeeze out results on when a deficiency 1 group contains a non-abelian free group (or equivalently contains F_2). In [57] J. S. Wilson conjectures that if a finitely presented group G is such that $\text{def}(G) + d^2/4 - d > 0$, where d is the minimum number of generators for G/G' (and assumed to be at least 2), then G contains a non-abelian free group. (Bartholdi has recently proved in [1] that such a group is non-amenable.)

For $\text{def}(G) = 1$ this was established in [39], as Corollary 2.4 in that paper states that if G is of deficiency 1 and $N \trianglelefteq G$ is such that G/N is non-trivial and free abelian with $N/N' \otimes_{\mathbb{Z}} \mathbb{F}$ having dimension at least 2 for some field \mathbb{F} , then G contains F_2 . Therefore on putting $N = \ker \chi$ for any χ , we have that $\beta_1(G; \mathbb{Z}/p\mathbb{Z})$ is at least 3 for some prime p which implies that $\beta_1(N; \mathbb{Z}/p\mathbb{Z}) \geq 2$.

We also have

Corollary 3.4 *If G has deficiency 1 and virtual first Betti number at least 2 then G contains F_2 or is $\mathbb{Z} \times \mathbb{Z}$ or the Klein bottle group.*

Proof. By [11] Theorem D, if a finitely presented group G does not contain F_2 but $\beta_1(G) \geq 2$ then there exists $K = \ker \chi$ which is finitely generated and such that $G/K \cong \mathbb{Z}$. Consequently by [41] Theorem 3 mentioned above, we have that K is of type FP_2 and hence free as before. Thus we are done unless K is free of rank 0 (but then $G = \mathbb{Z}$ so its virtual first Betti number is 1) or free of rank 1, giving the two exceptions. □

In fact it is widely believed that if G has deficiency 1 then it contains F_2 unless $G = BS(1, n)$. It is clear that the only case left is when $\beta_1(G) = 1$ with $1 \leq d(G/G') \leq 2$ and G is a strictly ascending HNN extension, so that S^0 contains two points with Σ one of them. The conjecture at the end of Section 2 in [10] is that in this case G is a strictly ascending HNN extension with base a finitely generated free group. As it is proved in this paper that for G a finitely presented group $\Sigma = \Sigma^1(G; \mathbb{Z})$ is equal to the higher dimensional invariant $\Sigma^2(G; \mathbb{Z})$, we would be done if we knew that $[\chi] \in \Sigma^2(G; \mathbb{Z})$ for a rationally defined χ implies that G is an ascending HNN extension over a base group of type FP_2 with associated homomorphism χ , just as is the case for $[\chi] \in \Sigma$ and with base of type FP_1 , i.e. finitely generated. In the special case of 2-generator 1-relator groups the conjecture is known: indeed

in [47] Chapter II Section 5 we have a “Tits alternative” which says that a subgroup of a 1-relator group either contains F_2 or is soluble (whereupon it is locally cyclic, $BS(1, n)$ or infinite dihedral). However we will not get such an alternative for arbitrary subgroups of deficiency 1 groups, or even arbitrary finitely generated subgroups:

Example 3.5

Thompson’s group T has a finite presentation of two generators and two relators, but has the unusual property that it does not contain F_2 nor is it virtually soluble. The group $G = T * \mathbb{Z}$ has deficiency at least one, and in fact it is exactly 1 as we can use Philip Hall’s inequality that $\text{def}(G) \leq \beta_1(G) - d(H_2(G; \mathbb{Z}))$ (see [55] 14.1.5). We have $H_n(T; \mathbb{Z}) = \mathbb{Z} \times \mathbb{Z}$ for all $n \geq 1$ by [18] Theorem 7.1, so that by using the Meier-Vietoris sequence for a free product we get $\beta_1(G) = 3$ and $H_2(G; \mathbb{Z}) = \mathbb{Z} \times \mathbb{Z}$.

Therefore any formulation of a possible Tits alternative for subgroups of deficiency 1 groups will need to avoid freely decomposable examples.

4 2-generator 1-relator groups of height 1

So far our attempts to prove that various deficiency 1 groups are large have needed some hypothesis on the group, such as being free-by-cyclic or LERF. In this section we concentrate on 2-generator 1-relator groups and look for conditions where we can conclude that such a group is large using information obtained directly from a given presentation. We fall into two very different cases: a group G of the form $\langle x, y | r(x, y) \rangle$ has $\beta_1(G)$ equal to either 1 or 2, the latter occurring exactly when $r \in F_2'$. First let us concentrate on the former case, where by making a change of free basis for the group F_2 we can assume that the presentation is of the form $\langle a, t | w(a, t) \rangle$ where the word w has exponent sum 0 in t and is cyclically reduced. Consequently we can define the height of such words: we can rewrite w as a word in $a_i = t^i a t^{-i}$ for $i \in \mathbb{Z}$ but only finitely many letters a_i will actually appear in w (this is sometimes referred to as Moldavanskii rewriting). If a_m is the smallest and a_M the largest letter to appear then the height of w is defined to be $M - m \geq 0$, which is invariant under cyclic permutations and taking inverses.

If the height is zero then we can only have $w = a^i$ so G is large (or \mathbb{Z} for $|i| = 1$). We now consider height one words where without loss of generality

we can assume that

$$w = ta^{i_1}t^{-1}a^{i_2} \dots ta^{i_{2k-1}}t^{-1}a^{i_{2k}}. \quad (2)$$

Note that for $\chi(t) = 1, \chi(a) = 0$ we have $\Delta_{G,\chi}(t) = (i_1 + \dots + i_{2k-1})t + i_2 + \dots + i_{2k}$ and for $\beta_1(G) = 1$ this is the Alexander polynomial Δ_G .

If $\beta_1(G) = 2$ then the exponent sum of x and y is zero, so we can talk about the height of either letter. Moreover a change of basis will preserve this property but could well vary the heights. The results that follow are also satisfied by groups G with $\beta_1(G) = 2$, where we interpret a height 1 word as there exists a free basis where the word has height 1 with respect to one of the letters.

Theorem 4.1 *If $G = \langle a, t | w \rangle$ where w is a height 1 word then either G is large or the finite residual $R_G = G''$, in which case G/R_G is metabelian and all finite images of G are metacyclic.*

Proof. By writing w in terms of a_0, a_1 we see that G is an HNN extension of the 1-relator group $H = \langle a_0, a_1 | w(a_0, a_1) \rangle$ with associated cyclic subgroups $A_0 = \langle a_0 \rangle$ and $A_1 = \langle a_1 \rangle$. We can now use Lubotzky's result as mentioned in the previous section: although it appears to require that G is LERF, all we need to conclude largeness for G is the existence of a homomorphism of G onto a finite group with $\theta(H) \neq \theta(A_0)$. But $\theta(A_0)$ is cyclic so if there is no such homomorphism then, as t conjugates a_0 into H , we have that $\theta(H)$ is normal in $\theta(G)$ and cyclic, with $\theta(G)/\theta(H) \cong \langle \theta(t) \rangle$ cyclic as well. This means that all finite images of G are metacyclic and if $x \in G''$ but $x \notin R_G$, we would take a homomorphism θ from G onto a finite group F with $\theta(x) \neq e$ so that F'' is non-trivial, but this is a contradiction. Consequently $G'' \leq R_G$ but Philip Hall's result that a finitely generated metabelian group is residually finite implies that $R_G \leq G''$. □

Moreover note that the normal closure of the element a in G/R_G , which is generated by elements of the form $t^i a t^{-i}$ for $i \in \mathbb{Z}$, is abelian because if there existed $[t^i a t^{-i}, a] \notin R_G$ then we could take a finite image of G in which this was also non-trivial, but this contradicts the fact that the cyclic group $\langle a \rangle$ is normal in every finite image.

This gives us a strict dichotomy in the behaviour of height 1 presentations. However the result does not tell us how to work out in which category a given presentation falls. Luckily there is a straightforward way of determining largeness once a major result of Zelmanov is used.

Corollary 4.2 *If $G = \langle a, t | w \rangle$ where w is a height 1 word then G is large if and only if there exists $H \leq_f G$ with $d(H/H') \geq 3$, where d is the minimum number of generators of a finitely generated group.*

Proof. The only if direction is clear. [58] states that if we have a finite presentation P of a group G with $\text{def}(P) + d^2/4 - d > 0$ where $d = d(G/G')$ is at least 2 then there exists a prime p such that the pro- p completion of G contains a non-abelian free pro- p group. In our case the deficiency is 1 so this is true if $d \geq 3$. But if $G'' = R_G$ then all finite images F of G have F'' trivial, so this is also true for the profinite or pro- p completion of G . Consequently any pro- p completion is soluble and cannot contain a non-abelian free group.

However G is 2-generated so we cannot have $d(G/G') \geq 3$ in any case. But for $H \leq_f G$ we still have deficiency 1, with $H'' \leq G''$ and $R_G = R_H$, so if $G'' = R_G$ and $d(H/H') \geq 3$ we have a contradiction. \square

We now give a summary of height 1 group presentations that have appeared in the literature. If we define the length of height 1 words to be k in (2) then it is clear that length 1 words are just the standard presentations of the Baumslag-Solitar groups $BS(m, n)$. In [4] Baumslag introduced a family $C(m, n)$ of groups with presentations

$$\langle a, t | (tat^{-1}a^m(tat^{-1})^{-1} = a^n) \rangle$$

which generalises the group $C(1, 2)$ that first appeared in [2] with the comment that they are 1-relator groups which are as far from being residually finite as possible. For our purposes we will interpret a finitely generated group G as being “as far from being residually finite as possible” in three ways: the first is that G (is infinite but) has no proper finite index subgroups whatsoever, or equivalently $G = R_G$. However this will never arise with deficiency 1 groups as G/G' is infinite.

The next strictest interpretation is that (G is non-abelian but) $G' = R_G$, or equivalently every finite image is abelian. To be succinct, we will say that G is **proabelian** because this is equivalent to the profinite completion of G being proabelian (or abelian). This property of finitely generated groups was examined in [24]. A finite index subgroup of a proabelian group is also proabelian and a finitely presented proabelian group G (with $\beta_1(G) > 0$) has $\Delta_G = 1$ (which can be seen because the Alexander polynomial of G is the same as that of G/R_G and this will be 1 for finitely generated infinite abelian groups). If G is a deficiency 1 group with the abelianisation G/G' equal

to $\mathbb{Z} \times T$ then G being proabelian and $\beta_1(G) = 1$ implies that T must be trivial because here $|\Delta_G(1)| = |T|$. We cannot have $d(G/G') \geq 3$ from above, so that if G is proabelian and $\beta_1(G) \geq 2$ then $G/G' = \mathbb{Z} \times \mathbb{Z}$. Moreover if $H/H' = \mathbb{Z}$ for all $H \leq_f G$ (or H/H' is $\mathbb{Z} \times \mathbb{Z}$ throughout) then G is proabelian by [24] Proposition 3.4. In particular if G has deficiency 1 with $\beta_1(G) = 2$ then either G is proabelian (so that $G/R_G = \mathbb{Z} \times \mathbb{Z}$) or a pro- p completion of G contains a non-abelian free pro- p group (so that G/R_G is “big”).

The first example of a proabelian finitely generated group that is given by a 1-relator presentation is $C(1, 2)$; indeed every finite image is cyclic as the abelianisation of $C(1, 2)$ is \mathbb{Z} .

Finally we could also look at when (G is not metabelian but) $G'' = R_G$, or equivalently every finite image is metabelian, for which we will similarly write that G is **prometabelian**. As we have $R_G \leq G''$ for any finitely generated group (although not necessarily $R_G \leq G'''$) this is also a natural concept, with a prometabelian group G being proabelian if and only if $G' = G''$. Moreover it is also fair to say that finitely generated proabelian and prometabelian groups are “as far from being large as possible”.

If we have a height 1 group $G = \langle a, t | w(a_0, a_1) \rangle$ which is not large then by the comment after Theorem 4.1 we have that a_0 commutes with a_1 in G/R_G . In particular if the exponent sum of a_0 in $w(a_0, a_1)$ is d and that of a_1 is c then G/G'' has the presentation

$$\langle a, t | ta^c t^{-1} = a^{-d}, [t^i a t^{-i}, a] \text{ for } i \in \mathbb{Z} \rangle$$

which is the same as for $BS(c, -d)/BS(c, -d)''$. Thus any result which depends purely on the finite images of $BS(m, n)$ for m and n coprime applies identically for any non-large group G of height 1 (and if c and d are not coprime then G is large anyway). In particular the formulae for the number of finite index subgroups of $BS(m, n)$ in [32] and the number of finite index normal subgroups in [23] apply equally for G . However there is a potential problem here: we do not know whether these groups are merely nonstandard presentations of $BS(m, n)$. For instance on putting $b = a^2$ and substituting in for the standard presentation of $BS(2, 3)$ we have $a = tbt^{-1}b^{-1}$ and so we obtain an alternative height 1 presentation

$$\langle b, t | tbt^{-1}b^{-1}tbt^{-1}b^{-2} \rangle.$$

The problem is a lack of invariants which are able to distinguish that two given presentations do not define isomorphic groups. We see in these cases

that any such invariant which is calculated using information obtained solely from the finite images of a group is here doomed to failure.

As for 1-relator proabelian groups that are not abelian, we can only have presentations with 2 generators. We have already mentioned $C(1, 2)$ above and in [29] the example $C(2, 3)$ was introduced where it was shown that for all $H \leq_f C(2, 3)$ we have $H/H' = \mathbb{Z}$ so $C(2, 3)$ is proabelian by [24]. Moreover if we regard $C(2, 3)$ as an HNN extension of $BS(2, 3)$ by adding the stable letter s with $sas^{-1} = t$ then we can iterate this process indefinitely and keep on obtaining new 1-relator height 1 group presentations. This HNN extension when applied to $C(1, 2)$ was also used in [25] to give an example of a higher dimensional knot whose infinite cyclic cover is not simply connected but which has no proper finite covers. The group $C(1, 2)$ (which is sometimes referred to as the Baumslag-Gersten group) also appears in [33] where it is shown that its Dehn function grows faster than every iterated exponential, and again in [19] where it is shown that it is isomorphic to

$$\langle a, t | (ta^{2^k}t^{-1})a(ta^{2^k}t^{-1})^{-1} = a^2 \rangle$$

for any $k \geq 0$. The paper [20] considers a more general family containing the groups $C(m, n)$ where the conjugating element tat^{-1} can be ta^kt^{-1} . These were used in [14] to give a pair of non-isomorphic 1-relator groups, each of which is a homomorphic image of the other. Also the examples

$$\langle a, t | (t^kat^{-k})a(t^kat^{-k})^{-1} = a^2 \rangle$$

of proabelian groups for $k \geq 1$ are mentioned in [50]: note this implies that a proabelian 1-relator group can have arbitrary height.

We can incorporate all of these examples in the next result. Although the proof is really that of Higman in [35] when giving the first example of an infinite finitely generated simple group, the aim here is to express it in as general terms as possible.

Theorem 4.3 *Suppose that g and h are elements of a finite group G with $h^kg^mh^{-k} = g^n$ for integers k, m, n where $|m - n| = 1$. Suppose further that g and h have equal orders. Then this order is always 1.*

Proof. If r is the order of g and h then by considering $h^{rk}g^{m^r}h^{-rk}$ which is equal to g^{n^r} , we must have r dividing $m^r - n^r$. So if $r \neq 1$ and p is the smallest prime dividing r , we have $(mn^{-1})^r \equiv 1 \pmod{p}$ where we can assume

without loss of generality that n and p are coprime. Thus the order of mn^{-1} divides r and $p - 1$ so p divides $m - n = \pm 1$.

□

Corollary 4.4 *Suppose $H = \langle a, t | w(a, t) \rangle$ where w has exponent sum 0 in t and is of height 1. Then the HNN extension $G = \langle H, s \rangle$, where $sas^{-1} = t$, is large if $H/H' \neq \mathbb{Z}$ and proabelian if $H/H' = \mathbb{Z}$ but H is not large.*

Proof. $G = \langle a, s | w(a, sas^{-1}) \rangle$ is also a 1-relator group of height 1 and if the Alexander polynomial of H (with respect to the homomorphism $\chi(t) = 1, \chi(a) = 0$) is $f(t) = ct + d$ then that for G is $c + d$. Thus G is large by Howie's result as the mod p Alexander polynomial is zero unless $|c + d| = 1$. But $|f(1)|$ is the order of T , where $H/H' = \mathbb{Z} \times T$.

If $|T| = 1$ and H is not large then suppose the finite group $F = \theta(G)$. We have that in F the relation $ta^c t^{-1} = a^{-d}$ holds because $\theta(H)$ is a finite image of H and $H'' = R_H$. But t and a are conjugate in F via the image of s , so we can use Theorem 4.3 to conclude that $\theta(t) = \theta(a) = e$.

□

Note that this means the group $C(m, n)$ above is not large if and only if $|m - n| = 1$. This is in disagreement with [4] Theorem 4 where part 2 states (but proofs are not given) that $C(m, n)$ is proabelian if m and n are distinct primes. However in Section 3 of that paper it is mentioned that if $m = n + 1$ then $C(m, n)$ is proabelian so we believe that is what was meant in the theorem.

Corollary 4.2 would allow us, using a computer, to gather statistics on what proportions of height 1 groups G are large by searching through finite index subgroups H of G until $d(H/H') \geq 3$ (or until we give up on G), thus giving a lower bound for the number of large groups. We propose to return to this but content ourselves for now by noting that the computer tells us that the group $\langle a, t | ta^2 t^{-1} a^{-1} ta^{-1} t^{-1} a^{-1} \rangle$ is large but has Alexander polynomial $t - 2$ which is the same as $BS(1, 2)$, and forming the HNN extension of this group using $sas^{-1} = t$ results in a height 1 group which is large but with Alexander polynomial equal to 1.

Corollary 4.5 *If $G_0 = \langle a, t | w \rangle$ is proabelian then $G_1 = \langle a, t | v^k w^m v^{-k} = w^n \rangle$ is proabelian, where v is a conjugate element of w in F_2 and $|m - n| = 1$.*

Proof. Given any finite image F of G_1 , we set h equal to the image of v and g equal to the image of w and apply Theorem 4.3. We conclude that the

image of w is trivial in F so F is a finite image of G_0 and hence is abelian. \square

Thus we see how to create lots of proabelian 1-relator groups by starting with $w = a$ (or indeed w equal to any element of a free basis for F_2) and then iterating in Corollary 4.5. We can also give an example of a 1-relator group G with $\beta_1(G) = 2$ that is proabelian, the existence of which was not previously known. Indeed the only known example up to now of such a group G which is not residually finite was given by Ol'shanskii in [52] but it is not directly constructive.

Corollary 4.6 *The group*

$$G = \langle a, t | [a, t][t, a^{-1}][a, t]^{-1} = [t, a^{-1}]^2 \rangle$$

is proabelian (hence not large) with $\beta_1(G) = 2$ but is not equal to $\mathbb{Z} \times \mathbb{Z}$ (hence is not residually finite).

Proof. On applying Corollary 4.5 with $w = [t, a^{-1}]$, so that $G_0 = \mathbb{Z} \times \mathbb{Z}$ and $v = [a, t]$, we obtain the above presentation for $G = G_1$. But it is known by [48] Section 4.4 that a 2-generator 1-relator group $\langle a, t | r \rangle$ is equal to $\mathbb{Z} \times \mathbb{Z}$ if and only if $r = [a, t]^{\pm 1}$ or a cyclic conjugate when r is cyclically reduced. \square

We note that the relation above is of height 1 with respect to t . However there are also groups with first Betti number 2 and height 1 which we can prove are large:

Corollary 4.7 *If $G = \langle a, t | w(a, t) \rangle$ where $\beta_1(G) = 2$ and w is a height 1 word of the form in (2) then G is large if $|i_1 + i_3 + \dots + i_{2k-1}| \neq \pm 1$ and is proabelian or large otherwise.*

Proof. We have that $\beta_1(G) = 2$ implies $i_1 + i_2 + \dots + i_{2k} = 0$ and so $\Delta_{G, \chi}(t) = n(t - 1)$ where $n = i_1 + i_3 + \dots + i_{2k-1}$ with $\chi(a) = 0$, $\chi(t) = 1$. Thus if $|n| \neq \pm 1$ we have largeness because $\Delta_{G, \chi}^p \equiv 0$ modulo a suitable prime p . We also have largeness by Corollary 4.2 unless $H/H' = \mathbb{Z} \times \mathbb{Z}$ for all $H \leq_f G$ which implies that G is proabelian (or $G = \mathbb{Z} \times \mathbb{Z}$ if $w = ta^{\pm 1}t^{-1}a^{\mp 1}$). \square

For instance the group $\langle a, b, t | aba^2 = b^2, ta^3t^{-1} = b \rangle$ is shown in [15] Proposition 19 to be word hyperbolic and of the form $F_6 \rtimes_{\alpha} \mathbb{Z}$. As such groups are residually finite and eliminating b gives a height 1 word with respect to t , we can use Corollary 4.7 to obtain another group satisfying the conditions of Corollary 2.4 without using the computer.

5 Problem List

Here we list some problems in Group Theory with particular emphasis on finitely generated and finitely presented groups. The aim is to put together questions which may well have appeared somewhere in print but which are not to be found in the standard problem lists (for instance the Kourovka notebook [42], the New York list [52] and Mladen Bestvina's questions in Geometric Group Theory [6]). Also there is an emphasis on questions which can be stated using only group theoretic concepts, although it may well be that solutions require topological, geometric or other techniques. In some cases the credits refer to those from whom we first heard about the question; we apologise if this is not the original source. The order of appearance places what should be the most general questions first and then specialises until we reach the topics of this paper.

Finiteness questions

1. (J. S. Wilson [57]) Is there a finitely presented and residually finite group which is neither virtually soluble nor contains a non-abelian free group?

Notes: If we replace finitely presented with finitely generated then certainly a range of examples are known. If however we keep finitely presented and remove residually finite then the list becomes much shorter. We know of only four constructions: Thompson's group F , the Houghton groups as in [38], the finitely presented Grigorchuk group (which is an ascending HNN extension of the well known Grigorchuk group) and the non-amenable monsters (which are ascending HNN extensions of finitely generated infinite groups with finite exponent) in [54]. All of these are far from being residually finite. Of course if we weaken finitely presented to finitely generated but strengthen residually finite to linear then there are no examples by the Tits alternative.

A variation might be to strengthen the conditions in other ways (e.g. LERF, coherent, of type FP) with the aim of proving such an example does not exist.

The Prüfer rank of a group is the supremum of the minimum number of generators over all finitely generated subgroups.

2. If a group has infinite Prüfer rank then does it have an infinitely generated subgroup?

Notes: This question has interesting implications either way. We say that a group has max (or is Noetherian) if it and every subgroup is finitely gener-

ated. The question of whether a group with max is virtually polycyclic was settled in the negative by Ol'shanskii's construction of Tarski monsters, but these have finite Prüfer rank. Moreover this property (like max) is preserved by subgroups, quotients and extensions. Thus a counterexample in question 2 would be a genuinely new example of a group with max as it would not be obtained from known examples by these operations.

If however the answer is no then a result of Lubotzky and Mann in [46] says that a finitely generated, residually finite group with finite Prüfer rank is virtually soluble. Thus a residually finite group with max would be virtually polycyclic, giving a positive answer to question 31 in the first Kourovka notebook, credited to M.I. Kargapolov.

3. If G/N and N are both virtually soluble then is G ?

Notes: This is true and straightforward with virtually soluble replaced by soluble.

4. Is there a finitely presented group that is elementary amenable but not virtually soluble?

Notes: In [37] Hillman and Linnell give an infinitely generated example and then use it to obtain a finitely generated example.

Amenability and properties (T) and (τ)

5. Do all finitely generated (or finitely presented) infinite amenable groups have a proper finite index subgroup?

Notes: The rationals show this is not true for infinitely generated groups. We have the dichotomy that no infinite discrete group can have property (T) and be amenable. For property (τ) (always assumed here to be with respect to all normal finite index subgroups) we have by [44] Proposition 3.3.7 that a finitely generated amenable group with infinitely many finite index subgroups does not have (τ), but groups with finitely many finite index subgroups trivially have property (τ). If the answer is yes then we regain our dichotomy for finitely generated/presented groups.

An old question asks if there is a finitely generated infinite simple group which is amenable. This is equivalent to having a finitely generated infinite amenable group with no finite index subgroups, as we can quotient out by a maximal normal subgroup. Infinitely generated simple amenable groups certainly exist, for instance the union $\cup A_n$ of alternating groups. It has also been asked whether there are finitely presented infinite simple amenable

groups, but those wishing to establish their existence would need to answer the next question in the negative.

6. Does an infinite finitely presented simple group necessarily contain a non-abelian free group?

Notes: This is not true for infinite finitely generated simple groups but for all known finitely presented examples the answer seems to be yes.

7. (Lackenby) If a finitely presented group has zero virtual first Betti number then does it have property (τ) ?

Notes: The Grigorchuk group shows that this is not true for finitely generated groups as it is amenable with infinitely many finite index subgroups.

We recall that property (T) implies finite generation and zero virtual first Betti number, as well as implying property (τ) which also implies zero virtual first Betti number.

Subgroup Separability

8. (J.S. Wilson) Is there a finitely generated group which is not virtually polycyclic but where every subgroup is the intersection of finite index subgroups?

Notes: The paper [40] shows that there are no examples when the group is hyper-(abelian or finite) which covers being virtually soluble. As for a finitely presented example, this would be residually finite and would not contain F_2 , so would provide a yes answer to question 1.

9. (Long and Reid [43] Question 4.5) Is there a finitely generated infinite group which is LERF and has property (T) ?

10. Is there a finitely presented infinite group which is LERF and which has zero virtual first Betti number?

Notes: A finitely presented example for Question 9 (or just one with property (τ)) would of course answer this. However the Grigorchuk group is a finitely generated infinite LERF group which has zero virtual first Betti number.

Largeness

11. If a finitely generated or finitely presented group has the biggest possible subgroup growth, that is of strict type n^n , then is it large?

12. Is there an algorithm to determine whether or not the group given by a finite presentation has a proper finite index subgroup?

Notes: It is unknown whether there is an algorithm to detect largeness in finitely presented groups. There are partial algorithms which will terminate with the answer yes if a finitely presented group is large and which may return no or not terminate otherwise, see for instance [22]. If we have a complete algorithm for largeness then, as pointed out by D. Groves and I. Leary, we have yes to this question because a group G has proper finite index subgroups if and only if $G * G * G$ is large.

13. If a finitely presented group has infinite virtual first Betti number then must it be large? Does it contain F_2 ? Must it be not virtually soluble?

Notes: These seem interesting questions. The first point to make is that these are all false for finitely generated groups, as in [28] it is pointed out that the soluble and \mathbb{R} -linear group $\mathbb{Z} \wr \mathbb{Z}$ has infinite virtual first Betti number. Therefore one would expect to find counterexamples by taking a group surjecting to $\mathbb{Z} \wr \mathbb{Z}$. But Baumslag shows in [3] Chapter IV Theorem 7 that a finitely presented group surjecting to $\mathbb{Z} \wr \mathbb{Z}$ is large.

We can also ask about growth of first Betti numbers in finite covers.

14. If a finitely presented group G has a sequence of finite index subgroups H_n such that $\beta_1(H_n)/[G : H_n] \geq c > 0$ for all n then is G large?

Notes: If $H_n \trianglelefteq G$ with the sequence nested and such that $\cap H_n$ is trivial (thus implying that G is residually finite) then this limit exists and is the first L^2 -Betti number (at least if $G = \pi_1(X)$ for X a CW-complex of finite type). However groups of the form $F_n \rtimes_{\alpha} \mathbb{Z}$ have zero first L^2 -Betti number but any large example will have a sequence as in the question.

If the sequence (H_n) above is such that there is a surjective homomorphism $\chi : G \rightarrow \mathbb{Z}$ with $K = \ker \chi \leq H_n$ for all n then G is large. Indeed we only require that $\beta_1(H_n)$ is unbounded, or even $\beta_1(H_n; \mathbb{Z}/p\mathbb{Z})$ for some prime p . This is because

$$\beta_1(H_n; \mathbb{F}) \leq \beta_1(K; \mathbb{F}) + 1 \text{ for } \mathbb{F} = \mathbb{Q} \text{ or } \mathbb{Z}/p\mathbb{Z}$$

so if we have an unbounded sequence then $\Delta_{G, \chi}^{\mathbb{F}}$ is zero, with Howie's result implying that for all large k the subgroup KG^k has a surjection onto $C_p * C_p$ (or $\mathbb{Z} * \mathbb{Z} * \mathbb{Z}$ for $\mathbb{F} = \mathbb{Q}$). However the counterexample $\mathbb{Z} \wr \mathbb{Z}$ above

for finitely generated groups also has the first Betti number growing linearly in a sequence of subgroups containing $\ker \chi$.

One could also formulate variations of question 14 using first Betti numbers modulo a prime p . However question 13 is no longer true in this more general setting: consider the \mathbb{R} -linear Baumslag-Remeslennikov soluble group

$$G = \langle a, s, t \mid tat^{-1} = asas^{-1}, [a, sas^{-1}] = 1 = [s, t] \rangle.$$

By using the binomial theorem modulo any prime p , it is not hard to show that we have subgroups H_n with $\beta_1(H_n; \mathbb{Z}/p\mathbb{Z}) \geq 2 + 2p^{n-1}$ where $G' \trianglelefteq H_n \trianglelefteq G$ is the finite abelian cover corresponding to

$$\begin{aligned} s^{-p^{n-1}}(t^{p^n+p^{n-1}}) \text{ and } s^{3p^{n-1}} & \text{ for } p \equiv 1 \pmod{3}, \\ t^{p^n+p^{n-1}} \text{ and } s^{3p^{n-1}} & \text{ for } p \equiv 2 \pmod{3}, \\ t^{p^n+p^{n-1}} \text{ and } s^{8p^{n-1}} & \text{ for } p = 3. \end{aligned}$$

This is the fastest possible growth of $\beta_1(H; \mathbb{Z}/p\mathbb{Z})$ because Corollary 1.4 in [57] states that if G is finitely presented and soluble then there exists $\kappa > 0$ such that $d(H/H') \leq \kappa |G : H|^{1/2}$ for all $H \leq_f G$.

15. Fix an integer $k \leq 1$. Is there $f(k)$ such that whenever a group G has a presentation of deficiency k and $\beta_1(G) > f(k)$ then G is large?

This seems less and less likely as k gets smaller. It is of course true for $k \geq 2$.

16. (J.S. Wilson [57]) If a group G has a finite presentation P such that $\text{def}(P) + d^2/4 - d > 0$ where $d \geq 2$ is the minimum number of generators of G/G' then does G contain a non-abelian free group?

Notes: Bartholdi has recently proved in [1] that Golod-Shafarevich groups are amenable. These are groups with a presentation (of finitely many generators but possibly infinitely many relators) possessing a certain condition which is certainly satisfied by finite presentations of the form above. This can be seen as providing evidence for the answer yes because a counterexample would have the extremely rare property of being a finitely presented non-amenable group without F_2 as a subgroup. Lackenby and Lubotzky ask for a group with a finite presentation as above which is not large: surely these exist? There is also the question (credited to Lubotzky and Zelmanov) of whether a group with such a presentation can have property (T) or (τ) . Recently Ershov in [30] has constructed Golod-Shafarevich groups with property (T) .

Deficiency 1 Groups

17. Is every deficiency 1 group that does not contain F_2 isomorphic to $BS(1, m)$?

Notes: For evidence, see Section 3.

18. Is every residually finite deficiency 1 group that is not large isomorphic to $BS(1, m)$?

Notes: In particular are all word hyperbolic groups of the form $F_n \rtimes_{\alpha} \mathbb{Z}$ for $n \geq 2$ large? A proof could be attempted in two parts: assume the virtual first Betti number is at least 2 to obtain largeness, and separately establish this assumption which is question 12.16 in [6] due to Casson.

19. If a deficiency 1 group G has (virtual) first Betti number at least 3 then is it large? If the minimum number of generators of G/G' (or H/H' for $H \leq_f G$) is at least 3 then is G large?

Notes: The first part is question 15 for $k = 1$ and $f(1) = 2$ which is certainly more believable here. Corollary 4.6 shows that we cannot replace “(virtual) first Betti number at least 3” with “(virtual) first Betti number at least 2 and $G \neq \mathbb{Z} \times \mathbb{Z}$ (or $BS(1, -1)$)”, even in the 1-relator case.

2-generator 1-relator presentations

In this special case question 17 is true but questions 18 and 19 are unknown.

20. Let G be a group with a 2-generator 1-relator presentation where the relator is not a proper power. Suppose that G is residually finite then does G have a finite index subgroup H which is an ascending HNN extension of a finitely generated free group?

Notes: If so then every finite index subgroup of H , and G , has deficiency 1. However if the relator is a proper power, which is exactly when G has torsion, then often G has a finite index subgroup of deficiency at least 2 and in any case G is known to be large (although [4] Problem 4 asks if a 1-relator group with torsion is virtually free-by-cyclic: here the free part could be infinitely generated and being cyclic includes trivial here). But if G is torsion free then it has geometric dimension 2 so all finite index subgroups have deficiency 1.

If we have a yes answer to this question then by [31] H , and so G , is coherent which is a special case of the old question of whether all 1-relator groups are coherent.

21. (Borisov and Sapir [13]) Is the property of being residually finite generic amongst 2-generator 1-relator presentations?

Notes: It is shown in [27] Theorem 6.1 that a generic presentation is not an ascending HNN extension of a finitely generated free group, but experimentally 94% of presentations give groups of the form $F_n \rtimes \mathbb{Z}$. There is an algorithm, due to Moldavanskii [49] in the case $\beta_1(G) = 1$ and Brown [17] for $\beta_1(G) = 2$, as to whether a 2-generator 1-relator group is of this form. One could try looking for $H \leq_f G$ with H an ascending HNN extension of a finitely generated free group (which is shown to be residually finite in [13]) but the problem is of course that although H will have a deficiency 1 presentation, it will not in general be 2-generated (indeed as soon as $d(H/H') \geq 3$ it cannot be).

We can also ask about genericity for other properties: being linear, being coherent (as evidence for the general conjecture) and being large. It is known that a generic presentation gives rise to a group that is one ended, torsion free and word hyperbolic, so if $G = F_n \rtimes \mathbb{Z}$ is large for word hyperbolic G with $n \geq 2$ as in question 18 then we would expect at least 94% of random presentations to give large groups.

22. Given reduced words $u, v \in F_2 = \langle x, y \rangle$ which are not a generating pair, does there exist a homomorphism ρ from F_2 into the symmetric group Σ_n for some n such that $\rho(u)$ and $\rho(v)$ commute but $\rho(x)$ and $\rho(y)$ do not?

Notes: This is equivalent to asking: if the group $G = \langle x, y | [u, v] \rangle$ is proabelian then is it abelian (and thus equal to $\mathbb{Z} \times \mathbb{Z}$)? Again there are interesting consequences because if so then G is either large or $\mathbb{Z} \times \mathbb{Z}$ by [24] Theorem 3.6 but if not then [52] Question (OR8) is answered, which asks if G is always residually finite.

23. (Müller and Schlage-Puchta [51] Problem 5) Is there an algorithm to detect whether a 2-generator 1-relator presentation is large?

This special case of question 12 might be more likely to yield the answer yes. However even if there is a straightforward criterion for largeness of such presentations, it might not convert into an algorithm: for instance if question 19 is true then it will detect largeness (which we can do anyway for an arbitrary finitely presented group) but would fail to prove that a given presentation is not large. We believe that Section 4 provides evidence that there is not an obvious algorithm.

References

- [1] L.Bartholdi, On amenability of group algebras, II: graded algebras. Available at <http://arxiv.org/abs/math/0611709>
- [2] G.Baumslag, A non-cyclic one-relator group all of whose finite quotients are cyclic. *J. Austral. Math. Soc.* **10** (1969), 497–498.
- [3] G.Baumslag, *Topics in combinatorial group theory*, Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel 1993.
- [4] G.Baumslag, Some open problems. In *Summer School in Group Theory in Banff, 1996*, CRM Proc. Lecture Notes, 17, Amer. Math. Soc., Providence 1999, 1–9.
- [5] B.Baumslag and S.J.Pride, Groups with two more generators than relators. *J. London Math. Soc.* **17** (1978), 425–426.
- [6] M.Bestvina, Questions in Geometric Group Theory. Available at <http://www.math.utah.edu/~bestvina>
- [7] M.Bestvina and M.Feighn, A combination theorem for negatively curved groups. *J. Differential Geom.* **35** (1992), 85–101.
- [8] M.Bestvina and M.Feighn, Addendum and correction to “A combination theorem for negatively curved groups”. *J. Differential Geom.* **43** (1996), 783–788.
- [9] M.Bestvina and M.Handel, Train tracks and automorphisms of free groups. *Ann. of Math.* **135** (1992), 1–51.
- [10] R.Bieri, Deficiency and the geometric invariants of a group. *J. Pure Appl. Algebra*. **208** (2007), 951–959.
- [11] R.Bieri, W.D.Neumann and R.Strebel, A geometric invariant of discrete groups. *Invent. Math.* **90** (1987), 451–477.
- [12] R.Bieri and R.Strebel, Almost finitely presented soluble groups. *Comment. Math. Helv.* **53** (1978), 258–278.

- [13] A. Borisov and M. Sapir, Polynomial maps over finite fields and residual finiteness of mapping tori of group endomorphisms. *Invent. Math.* **160** (2005), 341–356.
- [14] A. V. Borshchev and D. I. Moldavanskii, On the isomorphism of some groups with one defining relation. *Math. Notes*. **79** (2006), 31–40.
- [15] N. Brady and J. Crisp, CAT(0) and CAT(-1) dimensions of torsion free hyperbolic groups. *Comment. Math. Helv.* **82** (2007), 61–85.
- [16] P. Brinkmann, Hyperbolic automorphisms of free groups. *Geom. Funct. Anal.* **10** (2000), 1071–1089.
- [17] K. S. Brown, Trees, valuations, and the Bieri-Neumann-Strebel invariant. *Invent. Math.* **90** (1987), 479–504.
- [18] K. S. Brown and R. Geoghegan, An infinite-dimensional torsion-free FP_∞ group. *Invent. Math.* **77** (1984), 367–381.
- [19] A. M. Brunner, A group with an infinite number of Nielsen inequivalent one-relator presentations. *J. Algebra*. **42** (1976), 81–84.
- [20] A. M. Brunner, On a class of one-relator groups. *Canad. J. Math.* **32** (1980), 414–420.
- [21] J. O. Button, Mapping tori with first Betti number at least two. *J. Math. Soc. Japan*. **59** (2007), 351–370.
- [22] J. O. Button, Large mapping tori of free group endomorphisms. Available at <http://arxiv.org/abs/math/0511715>
- [23] J. O. Button, A formula for the normal subgroup growth of Baumslag-Solitar groups. *J. Group Theory* (to appear).
- [24] J. O. Button, Large groups of deficiency 1. *Israel J. Math.* (to appear).
- [25] J. O. Button, Finite covers of the infinite cyclic cover of a knot. *J. Knot Theory Ramifications* (to appear).
- [26] N. M. Dunfield, Alexander and Thurston norms of fibered 3-manifolds. *Pacific J. Math.* **200** (2001), 43–58.

- [27] N. M. Dunfield and D. P. Thurston, A random tunnel number one 3-manifold does not fiber over the circle. *Geom. Topol.* **10** (2006), 2431–2499.
- [28] M. Edjvet, The concept of “Largeness” in Group Theory. Ph.D. thesis, University of Glasgow, Glasgow 1984.
- [29] M. Edjvet and S. J. Pride, The concept of “largeness” in group theory II. In *Groups – Korea 1983*, Lecture Notes in Math. **1098**, Springer, Berlin 1984, 29–54.
- [30] M. Ershov, Golod-Shafarevich groups with property (T) and Kac-Moody groups. Available at <http://www.math.uchicago.edu/~ershov/Research/>
- [31] M. Feighn and M. Handel, Mapping tori of free group automorphisms are coherent. *Ann. of Math.* **149** (1999), 1061–1077.
- [32] E. Gelman, Subgroup growth of Baumslag-Solitar groups. *J. Group Theory*. **8** (2005), 801–806.
- [33] S. M. Gersten, Isoperimetric and isodiametric functions of finite presentations. In *Geometric group theory, Vol. 1 (Sussex, 1991)*, London Math. Soc. Lecture Note Ser. 181, Cambridge Univ. Press, Cambridge 1993, 79–96.
- [34] S. M. Gersten and J. R. Stallings, Irreducible outer automorphisms of a free group. *Proc. Amer. Math. Soc.* **111** (1991), 309–314.
- [35] G. Higman, A finitely generated infinite simple group. *J. London Math. Soc.* **26** (1951), 61–64.
- [36] J. A. Hillman, Tits alternatives and low dimensional topology. *J. Math. Soc. Japan*. **55** (2003), 365–383.
- [37] J. A. Hillman and P. A. Linnell, Elementary amenable groups of finite Hirsch length are locally-finite by virtually-soluble. *J. Austral. Math. Soc. Ser. A*. **52** (1992), 237–241.
- [38] C. H. Houghton, The first cohomology of a group with permutation module coefficients. *Arch. Math. (Basel)*. **31** (1978/79), 254–258.

- [39] J. Howie, Free subgroups in groups of small deficiency. *J. Group Theory*. **1** (1998), 95–112.
- [40] S. C. Jeanes and J. S. Wilson, On finitely generated groups with many profinite-closed subgroups. *Arch. Math. (Basel)*. **31** (1978/79), 120–122.
- [41] D. H. Kochloukova, Some Novikov rings that are von Neumann finite and knot-like groups. *Comment. Math. Helv.*. **81** (2006), 931–943.
- [42] *The Kourovka notebook. Unsolved problems in group theory*, Sixteenth edition, Edited by V. D. Mazurov and E. I. Khukhro. Russian Academy of Sciences Siberian Division, Institute of Mathematics, Novosibirsk 2006.
- [43] D. D. Long and A. W. Reid, Subgroup separability and virtual retractions of groups. Available at <http://www.math.utexas.edu/users/areid/publications.html>
- [44] A. Lubotzky, *Discrete groups, expanding graphs and invariant measures*. Progress in Mathematics 125, Birkhäuser Verlag, Basel 1994.
- [45] A. Lubotzky, Free Quotients and the first Betti number of some hyperbolic manifolds. *Transform. Groups*. **1** (1996), 71–82.
- [46] A. Lubotzky and A. Mann, Residually finite groups of finite rank. *Math. Proc. Cambridge Philos. Soc.*. **106** (1989), 385–388.
- [47] R. C. Lyndon and P. E. Schupp, *Combinatorial Group Theory*. Springer-Verlag, Berlin-Heidelberg-New York 1977.
- [48] W. Magnus, A. Karrass and D. Solitar, *Combinatorial Group Theory*, Dover Publications, Inc., Mineola, New York 2004.
- [49] D. Moldavanskiĭ, Certain subgroups of groups with one defining relation. *Sibirsk. Mat. Ž.* **8** (1967), 1370–1384.
- [50] D. Moldavanskiĭ and N. Sibyakova, On the finite images of some one-relator groups. *Proc. Amer. Math. Soc.*. **123** (1995), 2017–2020.
- [51] T. W. Müller and J.-C. Schlage-Puchta, Some examples in the theory of subgroup growth. *Monatsh. Math.*. **146** (2005), 49–76.

- [52] New York Group Theory Cooperative, Open problems in combinatorial and geometric group theory. Available at <http://zebra.sci.ccny.cuny.edu/web/nygtc/problems/>
- [53] A. Yu. Ol'shanskiĭ, SQ-universality of hyperbolic groups. *Sb. Math.* **186** (1995), 1199–1211.
- [54] A. Yu. Ol'shanskiĭ and M. V. Sapir, Non-amenable finitely presented torsion-by-cyclic groups. *Publ. Math. Inst. Hautes Études Sci.* **96** (2003), 43–169.
- [55] D. J. S. Robinson, *A course in the theory of groups*, Second edition, Graduate Texts in Mathematics 80, Springer-Verlag, New York 1996.
- [56] J. R. Stallings, Topologically unrealizable automorphisms of free groups. *Proc. Amer. Math. Soc.* **84** (1982), 21–24.
- [57] J. S. Wilson, Finitely presented soluble groups. In *Geometry and cohomology in group theory (Durham, 1994)*, London Math. Soc. Lecture Note Ser. 252, Cambridge Univ. Press, Cambridge 1998, 296–316.
- [58] E. Zelmanov, On groups satisfying the Golod-Shafarevich condition. In *New horizons in pro-p groups*, Progr. Math. 184, Birkhäuser Boston, Boston 2000, 223–232.